

Biceps Tenodesis Protocol

The intent of this protocol is to provide the therapist with guidelines of the post-operative rehabilitation course. It should not be a substitute for one's clinical decision making regarding the progression of a patient's post-operative course based on their physical exam findings, individual progress, and/or the presence of post-operative complications. The therapist should consult the referring physician with any questions or concerns.

Progression to the next phase based on clinical criteria and/or time frames as appropriate.

Phase I – Passive Range of Motion (Weeks 1- 2)

Goals

- Minimize shoulder pain and inflammatory response
- Achieve gradual restoration of passive range of motion (PROM)
- Enhance/ensure adequate scapular function

Precautions/Patient Education

- No active range of motion (AROM) of the elbow
- No excessive external rotation range of motion (ROM) / stretching. Stop when you feel the first end feel
- Use of a sling to minimize activity of biceps
- Ace wrap upper forearm as needed for swelling control
- No lifting of objects with operative shoulder
- Keep incisions clean and dry
- No friction massage to the proximal biceps tendon / tenodesis site
- Patient education regarding limited use of upper extremity despite the potential lack of or minimal pain or other symptoms

Activity

- Shoulder pendulum hang exercise
- PROM elbow flexion/extension and forearm supination/pronation
- AROM wrist/hand
- Begin shoulder PROM all planes to tolerance. Do not force any painful motion
- Scapular retraction and clock exercises for scapula mobility progressed to

scapular isometric exercises

- Ball squeezes
- Sleep with sling as needed supporting operative shoulder, place a towel under the elbow to prevent shoulder hyperextension
- Frequent cryotherapy for pain and inflammation
- Patient education regarding postural awareness, joint protection, positioning, hygiene, etc.
- May return to computer based work

Milestones to progress to Phase II

- Appropriate healing of the surgical incision
- Full PROM of shoulder and elbow
- Completion of phase I activities without pain or difficulty

Phase II – Active Range of Motion (Week 3)

Goals

- Minimize shoulder pain and inflammatory response
- Achieve gradual restoration of AROM
- Begin light waist level functional activities
- Wean out of sling by the end of the 2-3 postoperative week
- Return to light computer work

Precautions

- No lifting with affected upper extremity
- No friction massage to the proximal biceps tendon / tenodesis site

Activity

- Begin gentle scar massage and use of scar pad for anterior axillary incision
- Progress shoulder PROM to active assisted range of motion (AAROM) and AROM all planes to tolerance
- Lawn chair progression for shoulder
- Active elbow flexion/extension and forearm supination/pronation (No resistance)
- Glenohumeral, scapulothoracic, and trunk joint mobilizations as indicated (Grade I - IV) when ROM is significantly less than expected. Mobilizations should be done in directions of limited motion and only until adequate ROM is gained
- Begin incorporating posterior capsular stretching as indicated
- Cross body adduction stretch
- Side lying internal rotation stretch (sleeper stretch)
- Continued Cryotherapy for pain and inflammation
- Continued patient education: posture, joint protection, positioning, hygiene, etc.

Milestones to progress to Phase III

- Restore full AROM of shoulder and elbow
- Appropriate scapular posture at rest and dynamic scapular control with ROM and

functional activities

- Completion of phase II activities without pain or difficulty

Phase III - Strengthening Phase (Weeks 6-8)

Goals

- Normalize strength, endurance, neuromuscular control
- Return to chest level full functional activities

Precautions

- Do not perform strengthening or functional activities in a given plane until the patient has near full ROM and strength in that plane of movement
- Patient education regarding a gradual increase to shoulder activities

Activity

- Continue A/PROM of shoulder and elbow as needed/indicated
- Initiate biceps curls with light resistance, progress as tolerated
- Initiate resisted supination/pronation
- Begin rhythmic stabilization drills
- External rotation (ER) / Internal Rotation (IR) in the scapular plane
- Flexion/extension and abduction/adduction at various angles of elevation
- Initiate balanced strengthening program
 - Initially in low dynamic positions
 - Gain muscular endurance with high reps of 30-50, low resistance 1-3lbs)
 - Exercises should be progressive in terms of muscle demand / intensity, shoulder elevation, and stress on the anterior joint capsule
 - Nearly full elevation in the scapula plane should be achieved before beginning elevation in other planes
 - All activities should be pain free and without compensatory patterns
 - Exercises should consist of both open and closed chain activities
 - No heavy lifting should be performed at this time
- Initiate full can scapular plane raises with good mechanics
- Initiate ER strengthening using exercise tubing at 30° of abduction (use towel roll)
- Initiate side lying ER with towel roll
- Initiate manual resistance ER supine in scapular plane (light resistance)
- Initiate prone rowing at 30/45/90 degrees of abduction to neutral arm position
- Begin subscapularis strengthening to focus on both upper and lower segments
- Push up plus (wall, counter, knees on the floor, floor)
- Cross body diagonals with resistive tubing IR resistive band (0, 45, 90 degrees of abduction Forward punch
- Continued cryotherapy for pain and inflammation as needed

Milestones to progress to Phase IV

- Appropriate rotator cuff and scapular muscular performance for chest level activity
- Completion of phase III activities without pain or difficulty

Phase IV – Advanced Strengthening (Week 10+)

Goals

- Continue stretching and PROM as needed/indicated
- Maintain full non-painful AROM
- Return to full strenuous work activities
- Return to full recreational activities

Precautions

- Avoid excessive anterior capsule stress
- With weight lifting, avoid military press and wide grip bench press

Activity

- Continue all exercises listed above
 - Progress isotonic strengthening if patient demonstrates no compensatory strategies, is not painful, and has no residual soreness
- Strengthening overhead if ROM and strength below 90 degree elevation is good
- Continue shoulder stretching and strengthening at least four times per week
- Progressive return to upper extremity weight lifting program emphasizing the larger, primary upper extremity muscles (deltoid, latissimus dorsi, pectoralis major)
 - Start with relatively light weight and high repetitions (15-25)
- May initiate pre-injury level activities/vigorous sports if appropriate/cleared by MD

Milestones to return to overhead work and sport activities

- Clearance from MD
- No complaints of pain
- Adequate ROM, strength and endurance of rotator cuff and scapular musculature for task completion
- Compliance with continued home exercise program