

SLAP Repair:

The intent of this protocol is to provide the therapist with guidelines of the post-operative rehabilitation course after a SLAP Repair. It should not be a substitute for one's clinical decision making regarding the progression of a patient's post-operative course based on their physical exam findings, individual progress, and/or the presence of post-operative complications. The therapist should consult the referring physician with any questions or concerns.

INDIVIDUAL CONSIDERATIONS:

Phase I (Weeks 0-6)**Goals**

- Control inflammation and pain
- Use cryocuff 3-4 times per day
- Start PROM

Sling

- Wear sling continuously (except for therapy and showering) for 6 weeks

Therapeutic Exercises

- PROM of elbow and wrist
- Ball squeezes
- PROM of shoulder
- 90 degrees forward flexion in scapular plane
- 0 degrees of external rotation
- IR to posterior belt line

Phase II (Weeks 6-8)**Criteria**

- 90 degrees of passive forward flexion

Goals

- Full, painless PROM/AROM in forward flexion and internal rotation
- 30 degrees of external rotation

Sling

- Use sling for 4 weeks continuously except for shower and therapy
- May discontinue at 4-6 weeks

Therapeutic Exercises

- Continue to increase PROM and begin AROM to achieve full forward flexion
- External rotation at the side to 30 degrees maximum
- Full internal rotation

Phase III (Weeks 8-12)

Criteria

- Full passive internal rotation and forward flexion

Goals

- External rotation to 45 degrees
- AROM to PROM parameters

Therapeutic Exercise

- PROM/AROM
 - Full forward flexion
 - Full internal rotation
 - 45 degrees of external rotation
- Begin posterior capsular stretching
 - Cross arm stretch
 - Side lying internal rotation
- Increase active/passive external rotation at the side to full by 8 weeks
- Initiate strengthening program - avoid elbow resistive exercises until 3 months PO
 - Initiate IR/ER at neutral with tubing
 - Initiate forward flexion, scaption, empty can
 - Initiate sideling ER and triceps strengthening
 - Prone abduction with external rotation
 - Shoulder shrugs with resistance
 - Shoulder retraction with resistance
 - Prone rows
 - May begin internal and external rotation in the 90/90 position

Phase V (Months 3-6)

Criteria

- Full, painless PROM and AROM in all planes

Goals

- Maintain full PROM and AROM
- Progress with strengthening
- Return to functional activities at 3 months
- Return to overhead sports and contact activities at 6 months

Therapeutic Exercises

- Continue with passive stretching as needed to maintain full PROM
- Progress with strengthening, increasing resistance and repetition
- May initiate bench press, military press, and lat pull-downs
- Initiate and advance light biceps strengthening
- Initiate light plyometric drills at 12-16 weeks
- Initiate a throwing program at 16-20 weeks
- Sports specific drills can start at 16-20 weeks

Return to Sports

- May return to sports at 6 months when ROM is symmetric and painless
- Strength is 90% of contra-lateral side